


27

# GAP JUNCTIONS MEDIATE AGONIST-INDUCED OSCILLATORY CONTRACTIONS IN VASCULAR SMOOTH MUSCLE

S.W. Watts, M-L. Tsai, R. Loch-Caruso, R.C. Webb

Dept. of Physiology, Univ. of Michigan, Ann Arbor, MI 48109-0622

Uterine oscillatory contractions are perpetuated by gap junction complex formation [1]. We tested the hypothesis that agonist-induced oscillations in arteries [2,3] are also mediated by gap junctions. Helical strips of rat mesenteric and tail arteries were mounted in chambers for isometric force measurements. Tetraethylammonium (TEA; 1-30 mM) induced oscillatory contractions that were not endothelium-dependent. Oscillations were blocked by heptanol, a gap junction inhibitor (upper panel). Furthermore, in cultured mesenteric arterial cells, TEA increased Lucifer yellow dye transfer between neighbouring cells, a measure of junctional communication. Heptanol blocked dye transfer (lower panel). These data support a role for gap junction activity in vascular smooth muscle contraction.



## References:

- Loch-Caruso R, Juberg DR, Caldwell V, Corcos IA. Cultured myometrial cells establish communicating gap junctions. *Cell Biol Int Rep* 1990; 14: 905-915.
- Watts SW, Traub O, Lamb FS, Myers JH, Webb RC. Effect of ramipril of alpha-receptor mediated oscillatory contractions in tail artery of hypertensive rats. *Eur J Pharmacol* 1993; 242: 245-253.
- Lamb FS, Webb RC. Regenerative electrical activity and arterial contraction in hypertensive rats. *Hypertension*

28

# THE CONTRACTILE MECHANISMS OF SODIUM METAVANADATE IN ISOLATED RAT AORTA

Q. Zhou, N. Satake, S. Shibata

Department of Pharmacology, University of Hawaii, School of Medicine, Honolulu, Hawaii 96822, USA

Vanadate and vanadate-containing compounds play important roles in modulating various cellular functions. Structurally similar to phosphate, vanadate has dramatic effects on the activity of many enzymes, either being an inhibitor or a stimulator [1]. Vanadate has been reported to induce contraction of guinea pig aorta, rabbit colon, rat aorta, and canine saphenous vein [2,3]. However, the contractile mechanism of vanadate in smooth muscles has not been well understood. Therefore, in the present study, the mechanism of contractile effect of vanadate was investigated in rat aorta. Sodium metavanadate ( $\text{NaVO}_3$ ) ( $10^{-5}$  M -  $3 \times 10^{-3}$  M) induced contractile responses in a concentration-dependent manner. Removal of endothelium did not affect the response to  $\text{NaVO}_3$ .  $\text{NaVO}_3$  was the most potent agent to cause contraction as compared to vanadyl sulphate ( $10^{-5}$  M -  $3 \times 10^{-3}$  M) and vanadium trichloride ( $10^{-5}$  M -  $3 \times 10^{-3}$  M). The response to  $\text{NaVO}_3$  was inhibited by nifedipine ( $10^{-6}$  M), 2-nitro-4-carboxyphenyl-N,N-diphenylcarbamate (NCDC;  $3 \times 10^{-5}$  M), a phospholipase C inhibitor, and H-7 ( $10^{-5}$  M), a protein kinase C inhibitor. The response to  $\text{NaVO}_3$  was not inhibited by ouabain ( $10^{-3}$  M), prazosin ( $10^{-6}$  M), an  $\alpha_1$ -adrenoceptor inhibitor, methysergide ( $10^{-6}$  M), a serotonin receptor inhibitor, triptelennamine ( $10^{-6}$  M), a histamine receptor inhibitor, glyburide ( $10^{-6}$  M), a  $\text{K}_{\text{ATP}}$  channel blocker, apamin ( $10^{-6}$  M), a  $\text{K}_{\text{Ca}}$  channel blocker or  $\text{Mg}^{2+}$ -removal. In addition, the response to  $\text{NaVO}_3$  was inhibited by indomethacin ( $10^{-5}$  M), a cyclooxygenase inhibitor, and AA861 ( $10^{-5}$  M), a 5-lipoxygenase inhibitor. The response to arachidonic acid ( $10^{-6}$  M -  $3 \times 10^{-4}$  M) was also inhibited by indomethacin ( $10^{-5}$  M), AA861 ( $10^{-5}$  M), nifedipine ( $10^{-6}$  M) and NCDC ( $3 \times 10^{-5}$  M), but not by H-7 ( $10^{-5}$  M). Further, in the presence of indomethacin ( $10^{-5}$  M) and AA861 ( $10^{-5}$  M), H-7 ( $10^{-5}$  M) still inhibited the residual response to  $\text{NaVO}_3$ . In rat aorta,  $\text{NaVO}_3$  increased the levels of inositol monophosphate (IP) by about 20 times the basal level (control:  $147.4 \pm 19.96$  cpm/mg wet tissue) and prostaglandin  $\text{F}_{2\alpha}$  by about 2 times the basal level (control:  $88.8 \pm 13.6$  pg/mg wet tissue). Indomethacin ( $10^{-5}$  M), AA861 ( $10^{-5}$  M) and NCDC ( $3 \times 10^{-5}$  M) inhibited the IP increase. Indomethacin ( $10^{-5}$  M), but not AA861 ( $10^{-5}$  M), also inhibited the  $\text{PGF}_{2\alpha}$  increase. These results suggest that the response to  $\text{NaVO}_3$  is due to increased phosphoinositide metabolism and partly to subsequent increase in the metabolism of arachidonic acid.

## References:

- Voelkel NF, Czartolomna J. Vanadate potentiates hypoxic pulmonary vasoconstriction. *J Pharmacol Exp Ther* 1991; 259: 666-672.
- Shimada T, Shimamura K, Sunano S. Effects of sodium vanadate on various types of vascular smooth muscles. *Blood Vessels* 1986; 23: 113-124.
- Sunano S, Shimada T, Shimamura K, Moriyama K, Ichida S. Extra- and intracellular calcium in vanadate-induced contraction of vascular smooth muscle. *Heart Vessels* 1988; 4: 6-13.